

Anisotropy of Electrons Accelerated by a High-Intensity Laser Pulse

N.B. Narozhny*, M.S. Fofanov†

Moscow State Engineering Physics Institute, 115409 Moscow, Russia

(February 1, 2008)

We describe a realistic model for a focused high-intensity laser pulse in three dimensions. Relativistic dynamics of an electron submitted to such pulse is described by equations of motion with ponderomotive potential depending on a single free parameter in the problem, which we refer to as the "asymmetry parameter". It is shown that the asymmetry parameter can be chosen to provide quantitative agreement of the developed theory with experimental results of Malka *et al.* [Phys. Rev. Lett. **78**, 3314 (1997)] who detected angular asymmetry in the spatial pattern of electrons accelerated in vacuum by a high-intensity laser pulse.

41.75.Lx, 52.40.Nk, 52.75.Di

In their recent paper, Malka *et al.* [1] have reported experimental observation of electrons accelerated to relativistic energies by a high-intensity linearly polarized subpicosecond laser pulse in vacuum (see also comments on the paper [1] and the author's reply in Ref. [2–4]). This effect, known as high-intensity ponderomotive scattering, was discussed in detail in Ref. [5]. It occurs when the quiver amplitude imparted by the laser field to an electron becomes comparable to the focal spot radius of the laser beam. If the beam is Gaussian, the radial restoring force acting on the electron decays exponentially and the electron can be scattered out of the pulse.

The data of Malka *et al.* [1] show that the energies gained by the scattered electrons are in good quantitative agreement with calculations of electron trajectories in the polarization plane made with the first-order paraxial model for the laser field [1,3]. Nevertheless, the first order paraxial model predicts isotropic electron scattering [3] that is not supported by experimental results. Indeed, accelerated electrons were detected by Malka *et al.* only in the $(\mathbf{E}, \mathbf{k})$ plane, while no significant signal was detected after rotating the laser polarization direction by 90° [1]. In our opinion, this discrepancy between the theory and experiment is due to the following. In the first-order paraxial model focusing of a plane monochromatic wave only leads to an appearance of nonvanishing longitudinal components of electromagnetic fields in the focal region. However, the focusing is known to affect transverse components of the fields also (see, e.g. Ref. [6]). In particular [6], a plane monochromatic wave, polarized linearly along the x axis and propagating along the z axis, is converted by an aplanatic system to a converging spherical wave with non-vanishing y and x components of electric and magnetic fields, respectively.

In this Letter, we show that experimentally observed [1] anisotropy of ponderomotive electron scattering can be explained in the framework of a realistic model for the laser field developed in our recent paper [7]. Our model is based on an exact solution of Maxwell equations in three dimensions (3D), which can serve to describe a stationary, focused monochromatic laser beam with characteristic frequency ω and arbitrary intensity. Amplitudes of electric and magnetic fields in the model depend on radial coordinates as well as the coordinate along the direction of the beam propagation. These amplitudes are characterized by parameters R and $L = \omega R^2$, which can be interpreted as the focal spot radius and the Rayleigh length of the laser beam, respectively. The model admits different field configurations, which are determined by two coordinate functions satisfying certain second-order partial differential equations. Some special choice of these functions describes the Gaussian beams, which are widely used in optics. The model can be generalized by introducing temporal amplitude envelope $g(\varphi/\omega\tau)$, where φ is the relativistically invariant phase of the traveling wave, to describe a laser pulse with finite duration τ . (It is assumed that the function $g(\varphi/\omega\tau)$ is equal to unity at the point $\varphi = 0$ and decreases exponentially at the periphery of the pulse for $|\varphi| \gg \omega\tau$.) In this case the electric and magnetic fields of the model constitute an approximate solution of Maxwell equations with the second-order accuracy with respect to small parameters Δ and Δ' , defined as

$$\Delta' = 1/\omega\tau \lesssim \Delta = 1/\omega R \ll 1. \quad (1)$$

For a pulse propagating along the z direction, x and y components of the electric field oscillate with phase difference,

*E-mail: narozhny@theor.mephi.ru

†E-mail: fofanov@theor.mephi.ru

which depends on z and values of x and y coordinates of a point in the plane $z = \text{const}$. Moreover, the aforementioned phase difference depends also on φ . Therefore, one cannot ascribe some definite type of polarization to a tightly focused laser pulse. Nevertheless, for a weakly focused pulse ($\Delta \ll 1$), there always exists a region near the axis of the beam $r \ll R$, where the field properties are very close to those of a plane wave field. This region we call "the plane wave zone". It is reasonable to ascribe polarization of the field in this region to the beam as a whole. Hereafter we refer to the field of the pulse as linearly polarized in this sense only. For a tightly focused beam the focal spot radius is of the order of the wavelength and the plane wave zone doesn't exist. Therefore, only polarization of the parental beam incident on the focusing optical system can be ascribed to the focused beam in this case.

An arbitrary field, linearly polarized along the x axis, may be represented [7] as a superposition of E - and H -polarized waves (i.e. waves with the vectors $\vec{E}$ and $\vec{H}$ being perpendicular to the direction of the pulse propagation, compare with Ref. [8]). Relative contributions of E - and H -polarized waves to the resulting field are characterized by the "asymmetry parameter" μ

$$\mu = \frac{E_{x0}^h - E_{x0}^e}{E_{x0}^h + E_{x0}^e}, \quad -\infty < \mu < \infty, \quad (2)$$

where $E_{x0}^{e,h}$ are the x components of the electric field for E - and H -polarized waves at the focal point $\mathbf{r} = 0$ for $\varphi = 0$. Note, that in contrast to the amplitude, the quantities $E_{x0}^{e,h}$ can take both positive and negative values.

In the lowest approximation in Δ and Δ' , the averaged equations of motion of electrons (ponderomotive equations) in the field of the linearly polarized laser pulse take the form [7]

$$\begin{aligned} \frac{d\vec{q}_\perp}{d\varphi} &= -\Delta \frac{m}{q_-} \frac{\partial U}{\partial \vec{\rho}_\perp}, & \frac{d\vec{\rho}_\perp}{d\varphi} &= \Delta \frac{\vec{q}_\perp}{q_-}, \\ \frac{dq_-}{d\varphi} &= 0, & \frac{d\zeta}{d\varphi} &= \Delta^2 \frac{q_z}{q_-}. \end{aligned} \quad (3)$$

Here $q^\mu = \langle p^\mu \rangle$, where p^μ is the 4-momentum of the electron, $q_- = q_0 - q_z$, $\vec{\rho}_\perp = \langle \vec{r}/R \rangle$, $\zeta = \langle z/R \rangle$, brackets $\langle \rangle$ mean averaging over fast oscillations, and the ponderomotive potential U is defined by the expression

$$U = \frac{m\eta_0^2}{2} g^2(\varphi/\omega\tau) \left\{ |F_1|^2 + \mu^2 |F_2|^2 + \mu \cos 2\psi (F_1 F_2^* + F_1^* F_2) \right\}, \quad (4)$$

where $\tan \psi = \varrho_x/\varrho_y$, and η_0 is the value of the dimensionless field intensity parameter

$$\eta^2 = \frac{e^2 \langle \mathbf{E}^2 \rangle}{m^2 \omega^2}, \quad (5)$$

at the focal point at the moment $\varphi = 0$ ($\eta_0 = a/\sqrt{2}$, where a is the parameter of Malka *et al.*). Functions $F_i(\vec{\rho}_\perp, \zeta; \Delta)$ are chosen in the form corresponding to the Gaussian beam [7]

$$\begin{aligned} F_1 &= (1 + 2i\zeta)^{-2} \left\{ 1 - \frac{\varrho_\perp^2}{1 + 2i\zeta} \right\} \exp \left\{ -\frac{\varrho_\perp^2}{1 + 2i\zeta} \right\}, \\ F_2 &= -\varrho_\perp^2 (1 + 2i\zeta)^{-3} \exp \left\{ -\frac{\varrho_\perp^2}{1 + 2i\zeta} \right\}. \end{aligned} \quad (6)$$

The equation (4) shows that the ponderomotive potential U depends on the azimuthal angle ψ , and hence is generally speaking asymmetric. The potential U is symmetric only for the case $\mu = 0$. The shape of the ponderomotive potential in the plane $z = 0$ for $\varphi = 0$ is shown in Fig.1 for the cases $\mu = 0$ and $\mu = -1.55$. Figure 1a represents the ponderomotive potential for the standard case of Gaussian beam commonly used in literature, while Figure 1b illustrates the dramatic difference between the cases $\mu = 0$ and $\mu \neq 0$. For $\mu \neq 0$, the ponderomotive potential, possesses (besides the central peak) two extra maxima, which are located in the polarization plane. They arise as a result of the non-uniform intensity distribution in the plane $z = 0$ for $\mu \neq 0$. Locations of the additional maxima, as

well as their amplitudes, are determined by the value of μ . It is noteworthy, that the case $\mu = 0$ is the only one when E_y - and H_x -components of the electric and magnetic fields remain to be equal to zero outside the plane wave zone for the pulse polarized along the x axis.

We use Eqs. (3) and (4) for our analysis of free electron acceleration by a co-propagating intense laser pulse in vacuum under conditions close to those used in the experiments of Malka *et al.* [1]. The initial electron energy is taken to be $\varepsilon = 10$ keV ($v_0 = 0.2c$). The laser field parameters are: $\lambda = 1$ μm , $\eta_0 = 2.12$ (corresponds to the parameter $a = 3$ of Malka *et al.*), $R = 10$ μm and $\omega\tau = 480$. For the temporal envelope of the pulse $g(\varphi/\omega\tau)$ a sine-squared shape is taken. The asymmetry parameter is not determined experimentally and remains a free parameter of the problem. Its value, $\mu = -1.55$, has been chosen for better fitting of our computational results to the results of the experiment.

It is clear that the maximum energy will be gained by electrons that initially propagate exactly along the axis of the laser beam. However, the ponderomotive equations yield zero net energy transfer for such electrons, since they can "feel" the spatial gradient of the Gaussian laser field, and hence can be scattered out of the pulse, only due to the quiver motion which is absent for the average trajectory described by the ponderomotive equations. Nevertheless, there exist a family of trajectories with nonzero initial distances from the beam axis, for which the gained energy is close to its maximum value. The energy and scattering angle of the electrons depend also on the position at which the particle is overtaken by the pulse [1]. Indeed, the maximum energy is obtained for the electrons that experience the peak field of the laser and therefore meet the laser pulse at some distance before the focus. As a result, we obtain a 3D domain of injection positions of electrons obtaining the final kinetic energy $W \geq 0.9$ MeV at the scattering angle 39.5° .

Different cross-sections of this domain for the potential with $\mu = -1.55$ (Fig. 1b) are shown in Fig. 2. The longitudinal size of the domain is of the order of the Rayleigh length for the laser beam L , whereas its transverse size is much less than the focal spot radius R . The cross sections in Fig. 2 display high degree of radial anisotropy. Their shape, of course, is essentially determined by the type of the ponderomotive potential or by the value of μ . In particular, the domain of injection positions for the case (not presented here for the sake of compactness) of the potential shown in Fig. 1a with $\mu = 0$ is purely radial, in agreement with [3].

To obtain the angular distribution of scattered electrons we have calculated their trajectories numerically, applying the fourth-order Runge-Kutta method to the ponderomotive equations (3). Initial positions of the electrons were taken from the domain shown in Fig. 2. Since the shape of the cross-sections $z = \text{const}$ varies very slowly at intervals $\delta z \sim R$, we have considered equidistant planes $z = nR$ with $n = -27, -26, \dots, 5$. In each of these cross-sections, the electron injection positions were chosen randomly under condition that their density was constant and equal to $3 \times 10^{15}/R^2$. Physically, such procedure corresponds to uniformity of the initial electron beam. The total number of electron trajectories considered in such a way was more than 2×10^7 . We were interested only in those trajectories, which crossed the plane $z = 11.66$ cm at the points contained inside the ring with radii $r_1 = 8.99$ cm and $r_2 = 9.89$ cm. The latter conditions were determined by the position and angular size of the detector in the experiment [1].

The results of the calculations are presented in Fig. 3. We plot the normalized number of scattered electrons $\langle n \rangle$ with final energies $W \geq 0.9$ MeV as a function of the azimuthal scattering angle α . The values of $\langle n \rangle$ at any given α were obtained by averaging the number of scattered electrons over the range of α equal to the angular size of the detector used in the experiments [1]. It is easily seen, that the angular distribution of scattered electrons essentially depends on the parameter of asymmetry μ . At $\mu = 0$, which corresponds to the symmetrical Gaussian ponderomotive potential shown in Fig. 1a, the distribution is isotropic and purely radial (compare to the result of Ref. [3]). At the same time, if $\mu = -1.55$, the number of scattered electrons detected in the $(\mathbf{E}, \mathbf{k})$ plane ($\alpha = 0$) is about 30 times higher than that in the $(\mathbf{H}, \mathbf{k})$ plane ($\alpha = \pi/2$). This result is in good quantitative agreement with observations of Malka *et al.* [1].

The 30-fold anisotropy of accelerated electrons is clearly explained by asymmetry of the ponderomotive potential (4). The cross term in (4), besides asymmetric corrections to the radial force, gives rise to a tangential force which is responsible for pushing electrons out of the plain perpendicular to the polarization plane. The latter corresponds to minimum of the ponderomotive potential as function of azimuthal angle ψ , while the perpendicular plane to its maximum. Therefore one could be surprised that the number of electrons scattered at the angle $\alpha = \pi/2$ is not equal to zero. Certainly it is explained by complicated structure of the ponderomotive potential (4), namely by the fact that the cross term can change its sign at the periphery of the focus.

The asymmetry of the ponderomotive potential itself is determined by non-zero value of the parameter μ which characterizes relative contributions of E- and H-polarized waves to the resulting field. As far as we know, nobody has never controlled the parameter μ in experiments. The reason is evident. Before the work of Malka *et al* [1] there were no experiments where the three-dimensional intensity distribution of a laser pulse influenced physical results. Therefore even qualitative coincidence of our calculations with the results of the experiment [1] would give a deeper

insight into physics of the electron-laser interaction. However, it appeared that the model describes the experiment quantitatively. Of course, the quantitative agreement between the developed theory and the experiment is based on fitting of a single free parameter μ in the problem, which has not been measured experimentally. Therefore, from the standpoint of our model, the experiment of Malka *et al.* [1] could be considered as a probe for 3D field distribution in the laser pulse. The correctness of our approach could be verified by an independent experiment for another physical situation performed with the same laser system. Measurements of angular distribution of ATI electrons could serve as a good example of such experiment.

We thank M.V. Fedorov and V.D. Mur for fruitful discussions. This work was supported by the Russian Foundation for Basic Research under projects 00-02-16354 and 00-02-17078.

-
- [1] G. Malka, E. Lefebvre, and J.L. Miquel, Phys. Rev. Lett. **78**, 3314 (1997)
 - [2] K.T. McDonald, Phys. Rev. Lett. **80**, 1350 (1998)
 - [3] P. Mora and B. Quesnel, Phys. Rev. Lett. **80**, 1351 (1998)
 - [4] G. Malka, E. Lefebvre, and J.L. Miquel, Phys. Rev. Lett. **80**, 1352 (1998)
 - [5] F.V. Hartemann *et al.*, Phys. Rev. E **51**, 4833 (1995)
 - [6] A. Boivin and E. Wolf, Phys. Rev. **138**, B1561 (1965)
 - [7] N.B. Narozhny, M.S. Fofanov, JETP, **90**, 753 (2000)
 - [8] M. Born and E. Wolf, *Principles of Optics* (Pergamon Press, New York, 1964)

FIG. 1. Ponderomotive potential (4) in the plane $z = 0$ at $\varphi = 0$ for two values of asymmetry parameter: (a) $\mu = 0$ and (b) $\mu = -1.55$.

FIG. 2. Cross sections of 3D injection-position domain for electrons gaining the energy $W \geq 0.9$ MeV in the ponderomotive potential (4) with $\mu = -1.55$. z coordinates of the cross sections are (a) $z = -21R$, (b) $z = -16R$, (c) $z = -11R$, (d) $z = -6R$, (e) $z = -R$, and (f) $z = 2R$.

FIG. 3. Normalized number of scattered electrons with the energy $W \geq 0.9$ MeV as a function of azimuthal angle α for (a) $\mu = 0$ and (b) $\mu = -1.55$.





